

$\mathbf{F} = (2x - y)\mathbf{i} - yz^2\mathbf{j} - y^2z^2\mathbf{k}$, calculate $\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} ds$ using Stoke's theorem, where S is the half upper surface of the sphere $x^2 + y^2 + z^2 = 1$, $(0 \leq z \leq 1)$. [98 嘉大土木 5]

[解](1) 令 $f = x^2 + y^2 + z^2 \Rightarrow \mathbf{n} = \frac{2x\mathbf{i} + 2y\mathbf{j} + 2z\mathbf{k}}{\sqrt{(2x)^2 + (2y)^2 + (2z)^2}} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2x - y & -yz^2 & -y^2z^2 \end{vmatrix} = 2y(-z^2 + z)\mathbf{i} + \mathbf{k}$$

$$(\nabla \times \mathbf{F}) \cdot \mathbf{n} = 2xy(-z^2 + z) + z, \quad \mathbf{n} \cdot \mathbf{k} = z$$

$$\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} ds = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} \frac{dxdy}{|\mathbf{n} \cdot \mathbf{k}|} = \iint [2xy(-z^2 + z) + z] \frac{dxdy}{z}$$

其中 $2xy(-z^2 + z)$ 為 x (或 y) 的奇函數 $\Rightarrow \iint 2xy(-z^2 + z) \frac{dxdy}{z} = 0$

$$\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} ds = \iint dxdy = \pi \cdot 1^2 = \pi$$

(2) 由 Stokes 定理知 $\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} ds = \oint_C \mathbf{F} \cdot d\mathbf{r}$, where $C: x^2 + y^2 = 1$

令 $x = \cos \theta, y = \sin \theta$

$$\mathbf{F} \cdot d\mathbf{r} = (2\cos \theta - \sin \theta)\mathbf{i} + 0\mathbf{j} + 0\mathbf{k} \cdot (-\sin \theta\mathbf{i} + \cos \theta\mathbf{j})d\theta = (-\sin 2\theta + \sin^2 \theta)d\theta$$

$$\begin{aligned} \oint_C \mathbf{F} \cdot d\mathbf{r} &= \int_0^{2\pi} (-\sin 2\theta + \sin^2 \theta) d\theta = \int_0^{2\pi} \left(-\sin 2\theta + \frac{1 - \cos 2\theta}{2}\right) d\theta \\ &= \left(\frac{\cos 2\theta}{2} + \frac{\theta}{2} - \frac{\sin 2\theta}{4}\right) \Big|_0^{2\pi} = \pi \end{aligned}$$